## BREASTFEEDING IS BEST FOR BABIES

It is important that, in preparation for and during breastfeeding, mother eats a **neartiny**, **palanced dist**. Combined breast and bottle feeding in the first weeks of life may reduce the supply of mother own breast milk, and reversing the decision not to breastfeed is difficult. Always **consult healthcare professional for advice** about feeding your baby. If infant formula is used, manufacturer's instructions should be followed for use carefully.

## References:

Akman I, et al. Arch Dis Child, 91(5):417–9, 2006.

Benninga MA, et al. Gastroenterol, 150(6):1443–55, 2016

Bongers ME, et al. Nutr J, 6:8, 2007.

Brown MM, et al. Curr Opin Ophthalmol, 20(3):188–94, 2009.

lacono G, et al. Dig Liver Dis, 37(6): 432–8, 2005.

Indrio F, et al. Eur J Pediatr, 174(6):841-2, 2015.

Miller-Loncar C, et al. Arch Dis Child, 89(10):908–12, 2004.

Morris S. Econ Hist Rev, 54(3):525–45, 2001.

Partty A, et al. JAMA Pediatr, 167: 977–8, 2013.

Rautava P, et al. Pediatrics, 96(1):43-7, 1995. Sano F, et al. Eur J Clin Nutr, 60:1304-10, 2006.

Sano F, et al. Eur J Clin Nutr, 60:1304-10, 2006.

Schmelze H, et al. JPGN, 36:343-51, 2003.

Sommers T, et al. Am J Gastroenterol, 110(4):572-9, 2015.

Vandenplas Y, et al. J Pediatr Gastroenterol Nutr, 61(5): 531-7, 2015.

Vandenplas Y, *et al.* The 47th ESPGHAN Annual Meeting, 9–12 June, 2014, Jerusalem, Israel. Abstract #PO-N-0253.

Veitl V, et al. J Emahrüngsmed, 2:14-20, 2000.

Vik T, et al. Acta Paediatr, 98(8):1344-8, 2009.